

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
4 January 2001 (04.01.2001)

PCT

(10) International Publication Number
WO 01/00243 A1

- (51) International Patent Classification⁷: **A61K 47/48**
- (21) International Application Number: **PCT/IB00/00746**
- (22) International Filing Date: **2 June 2000 (02.06.2000)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:
9915231.6 29 June 1999 (29.06.1999) **GB**
- (71) Applicant (for GB only): **PFIZER LIMITED** [GB/GB];
Ramsgate Road, Sandwich, Kent CT13 9NJ (GB).
- (71) Applicant (for all designated States except GB, US):
PFIZER INC. [US/US]; 235 East 42nd Street, New York,
NY 10017 (US).
- (72) Inventor; and
- (75) Inventor/Applicant (for US only): **BILLOTTE, Anne**
[FR/GB]; Pfizer Central Research, Ramsgate Road, Sand-
wich, Kent CT13 9NJ (GB).
- (74) Agents: **WOOD, David, J. et al.**; Pfizer Limited, Rams-
gate Road, Sandwich, Kent CT13 9NJ (GB).
- (81) Designated States (*national*): AE, AL, AM, AT, AU, AZ,
BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK,
DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID,
IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT,
LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ,
PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT,
TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— With international search report.
- For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: **PHARMACEUTICAL COMPLEX**

(57) Abstract: This invention relates to a complex of eletriptan and a sulphobutylether-beta-cyclodextrin, or a pharmaceutically acceptable salt thereof, and to processes for the preparation of, pharmaceutical formulations including, and the used of, such a complex.



WO 01/00243 A1

PHARMACEUTICAL COMPLEX

This invention relates to a complex of eletriptan and a sulphobutylether-beta-cyclodextrin, or a pharmaceutically acceptable salt thereof, and to

5 processes for the preparation of, pharmaceutical formulations including, and the uses of, such a complex.

Eletriptan, 3-([1-methylpyrrolidin-2(R)-yl]methyl)-5-(2-phenylsulphonylethyl)-1H-indole, is disclosed in WO-A-92/06973. A preferred hydrobromide salt form of eletriptan is disclosed in WO-A-96/06842. WO-A-10 99/01135 discloses a pharmaceutical formulation comprising eletriptan hemisulphate and caffeine. WO-A-00/06161 describes the use of eletriptan for the prevention of migraine recurrence.

Eletriptan is a 5HT_{1B/1D} receptor agonist and has been shown to be highly effective for the treatment of migraine and the prevention of migraine 15 recurrence. Eletriptan has also been disclosed for the treatment of hypertension, emesis, depression, anxiety, an eating disorder, obesity, drug abuse, cluster headache, pain, chronic paroxysmal hemicrania and a headache associated with a vascular disorder.

In order to administer eletriptan, or a salt thereof, by the intranasal route, 20 it is desirable that a formulation does not produce unacceptable levels of effect, such as irritancy, on the nasal mucosae. It has been found that the formulations described in WO-A-99/01135 comprising eletriptan hemisulphate and caffeine are irritant on the nasal mucosae.

Accordingly, it is an object of the present invention to provide a well- 25 tolerated pharmaceutical formulation of eletriptan, or a salt thereof, that is suitable for administration by the intranasal route.


It is further object of this invention to provide a well-tolerated, stable, aqueous, pharmaceutical formulation containing eletriptan, or a salt thereof, that is suitable for parenteral, preferably, intranasal administration and which 30 enables the drug to have good bioavailability and rapid onset of action.

WO-A-91/11172 and WO-A-94/02518 disclose sulphoalkyl ether cyclodextrin derivatives.

WO-A-98/02186 discloses an inclusion complex of (a) an indole selective serotonin (5-HT_{1D}) agonist or a pharmaceutically acceptable salt thereof and (b)
 5 an unsubstituted or substituted beta- or gamma-cyclodextrin, together with pharmaceutical compositions thereof.

It has now been found that eletriptan, or a salt thereof, can form a complex with certain sulphobutylether-beta-cyclodextrin derivatives of the type disclosed in WO-A-91/11172. Although this complexation undesirably and
 10 unpredictably decreases the aqueous solubility of eletriptan, or a salt thereof, it unexpectedly and advantageously provides a complex which is well-tolerated when administered intranasally, principally since it has a negligible irritant effect on the nasal mucosae in comparison with both the known caffeine formulations of eletriptan disclosed in WO-A-99/01135 and other cyclodextrin complexes of
 15 eletriptan.

Accordingly, the present invention provides a complex of eletriptan and a cyclodextrin derivative of the formula (I):-



20

wherein

R^{1a-g}, R^{2a-g} and R^{3a-g} each independently represent -OH or -O(CH₂)₄SO₃H;

provided that at least one of R^{1a-g} represents -O(CH₂)₄SO₃H:
or a pharmaceutically acceptable salt thereof.

Pharmaceutically acceptable salts of particular interest associated with the eletriptan component are acid addition and base salts thereof. Suitable acid
5 addition salts are formed from acids which form non-toxic salts and examples are the hydrochloride, hydrobromide, hydroiodide, sulphate, bisulphate, nitrate, phosphate, hydrogen phosphate, acetate, maleate, fumarate, lactate, tartrate, citrate, gluconate, succinate, saccharate, benzoate, methanesulphonate, ethanesulphonate, benzenesulphonate, p-toluenesulphonate and pamoate
10 salts. Hydrobromide and sulphate, including hemisulphate, salts are preferred. Suitable base salts are formed from bases which form non-toxic salts and examples are the sodium, potassium, aluminium, calcium, magnesium, zinc and diethanolamine salts.

Pharmaceutically acceptable salts of particular interest associated with
15 the cyclodextrin ring component are base salts of the -O(CH₂)₄SO₃H groups, for example, alkali metal salts, such as sodium salts.

For a review on suitable salts see Berge *et al*, J. Pharm. Sci., **66**, 1-19, 1977.

A pharmaceutically acceptable salt may readily be prepared by mixing
20 together solutions of eletriptan, the cyclodextrin or the complex, and the desired acid or base, as appropriate. The salt may precipitate from solution and be collected by filtration or may be recovered by evaporation of the solvent. When a salt of eletriptan and/or the cyclodextrin are/is separately prepared, these/this may then be used for the preparation of the complex.

25 A polymorph of eletriptan, or a salt thereof, may also be used for the purpose of the present invention.

Preferably, the average number of -O(CH₂)₄SO₃H groups per molecule of formula (I) is in the range of from 6.1 to 6.9, for example, 6.5 or about 6.5.

It is preferred that each -O(CH₂)₄SO₃H present is in the form of an alkali
30 metal salt (such as the sodium salt).

Preferably, the molar ratio of eletriptan:cyclodextrin derivative of the formula (I) is from 1:1 to 15:1, most preferably from 1:1 to 10:1.

Preferably, eletriptan is present in the form of the hemisulphate salt.

The complex can be administered alone but will generally be

5 administered in admixture with a pharmaceutically acceptable excipient, diluent or carrier selected with regard to the intended route of administration and standard pharmaceutical practice.

For example, the complex can be administered orally, buccally or sublingually in the form of tablets, capsules, ovules, elixirs, solutions or
10 suspensions, which may contain flavouring or colouring agents, for immediate-, delayed-, sustained-, pulsed- or controlled-release applications.

Such tablets may contain excipients such as microcrystalline cellulose, lactose, sodium citrate, calcium carbonate, dibasic calcium phosphate and glycine, disintegrants such as starch (preferably corn, potato or tapioca starch),
15 sodium starch glycollate, croscarmellose sodium and certain complex silicates, and granulation binders such as polyvinylpyrrolidone, hydroxypropylmethylcellulose (HPMC), hydroxypropylcellulose (HPC), sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, stearic acid, glyceryl behenate and talc may be included.

20 Solid compositions of a similar type may also be employed as fillers in gelatin capsules. Preferred excipients in this regard include lactose, starch, a cellulose, milk sugar or a high molecular weight polyethylene glycol. For aqueous suspensions and/or elixirs, the complex may be combined with various sweetening or flavouring agents, colouring matter or dyes, with emulsifying
25 and/or suspending agents and with diluents such as water, ethanol, propylene glycol and glycerin, and combinations thereof.

The complex can also be administered parenterally, for example, intravenously, intra-arterially, intraperitoneally, intrathecally, intraventricularly, intrasternally, intracranially, intramuscularly or subcutaneously, or it may be
30 administered by infusion techniques. It is best used in the form of a sterile aqueous solution which may contain other substances, for example, enough

salts or glucose to make the solution isotonic with blood. The aqueous solutions should be suitably buffered (preferably to a pH of from 3 to 9), if necessary.

The preparation of suitable parenteral formulations under sterile conditions is readily accomplished by standard pharmaceutical techniques well-known to those skilled in the art.

For oral and parenteral administration to human patients, the daily dosage level of eletriptan will usually be from 0.001 to 0.50 mg/kg (in single or divided doses).

Thus tablets or capsules containing the complex may contain from 5 to 240 mg of eletriptan, or a salt thereof, for administration singly or two or more at a time, as appropriate. The physician in any event will determine the actual dosage which will be most suitable for any individual patient and it will vary with the age, weight and response of the particular patient. The above dosages are exemplary of the average case. There can, of course, be individual instances where higher or lower dosage ranges are merited and such are within the scope of this invention.

The complex can also be administered intranasally or by inhalation and is conveniently delivered as a single dose or multi-dose in the form of a dry powder inhaler or an aerosol spray presentation from a pressurised container, pump, atomiser, spray or nebuliser, with or without the use of a suitable propellant, e.g. dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, a hydrofluoroalkane such as 1,1,1,2-tetrafluoroethane (HFA 134A [trade mark]) or 1,1,1,2,3,3,3-heptafluoropropane (HFA 227EA [trade mark]), carbon dioxide or other suitable gas. In the case of a pressurised aerosol, the dosage unit may be determined by providing a valve to deliver a metered amount. The pressurised container, pump, atomiser, spray or nebuliser may contain a solution or suspension of the complex, e.g. using a mixture of ethanol and the propellant as the solvent, which may additionally contain a lubricant, e.g. sorbitan trioleate. Capsules and cartridges (made, for example, from gelatin) for use in an inhaler or insufflator

may be formulated to contain a powder mix of the complex and a suitable powder base such as lactose or starch.

Aerosol or dry powder formulations are preferably arranged so that each metered dose or "puff" contains from 1 to 16 mg of eletriptan for delivery to the
5 patient.

Alternatively, the complex can be administered in the form of a suppository or pessary, or it may be applied topically in the form of a lotion, solution, cream, ointment or dusting powder. The complex may also be transdermally administered, for example, by the use of a skin patch.

10 For application topically to the skin, the complex can be formulated as a suitable ointment containing the complex suspended or dissolved in, for example, a mixture with one or more of the following: mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene polyoxypropylene compound, emulsifying wax and water. Alternatively, it can
15 be formulated as a suitable lotion or cream, suspended or dissolved in, for example, a mixture of one or more of the following: mineral oil, sorbitan monostearate, a polyethylene glycol, liquid paraffin, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

The complex of the invention may be prepared from an aqueous solution
20 or slurry of eletriptan, or a salt thereof, and the cyclodextrin, or a salt thereof, by conventional methods. Where a solid complex is required, the solution or slurry may be dried by spray-drying or freeze-drying (lyophilisation). Alternatively, eletriptan, or a salt thereof, and the cyclodextrin, or a salt thereof, can be mixed and the powder mixture moistened with water. The mixture may then be
25 vigorously mixed to form a paste and dried at an elevated temperature, preferably under reduced pressure, to remove the water. The dried complex may be crushed and sieved to the desired particle size.

The solid complex can then be formulated for administration as a pharmaceutical formulation by standard methods, for example, if an aqueous
30 formulation is required the complex may be dissolved with stirring in water. A further suitable excipient, diluent or carrier may be incorporated in the

formulation at the blending or mixing step. Alternatively, a pharmaceutical formulation including the present complex may be prepared by mixing eletriptan, or a salt thereof, and the cyclodextrin, or a salt thereof, together with a suitable excipient, diluent or carrier directly.

5 The complex can also be administered together with a prokinetic (e.g. metoclopramide) or antiemetic agent.

It is to be appreciated that all references herein to treatment include curative, palliative and prophylactic treatment.

Preferably, the complex is administered as a pharmaceutical formulation,
10 most preferably, as an aqueous solution formulation.

Preferably, the pharmaceutical formulation is formulated for parenteral administration, for example, intranasal administration.

Preferably, a pharmaceutical formulation, most preferably, an aqueous pharmaceutical formulation, comprises one or more of the following excipients,
15 diluent or carriers:

- (a) an anti-oxidant, for example, citric acid or ascorbic acid;
- (b) a co-solvent, for example, ethanol, glycerol or a PEG200-400; and
- (c) an organic polymer, for example, a water-soluble organic polymer such as carboxymethylcellulose, polyvinylpyrrolidone or hydroxypropylmethyl
20 cellulose.

The presence of an anti-oxidant increases the stability of the formulation.

The presence of an organic polymer or co-solvent optimises the bioavailability and increases the absorption and the onset of action of eletriptan.

Preferably, the present invention provides a pharmaceutical formulation,
25 preferably an aqueous pharmaceutical formulation, including from 10 to 150 mg/g of eletriptan or a salt thereof (preferably the hemisulphate), and from 10 to 30% weight/weight of the sulphobutylether-beta-cyclodextrin.

Preferably, up to and including 1% weight/weight of an anti-oxidant such as citric acid or ascorbic acid can be present.

30 Preferably, up to and including 30% weight/weight of a co-solvent such as ethanol, glycerol or a PEG200-400, preferably glycerol, can be present.

Preferably, up to and including 0.5% weight/weight of an organic polymer such as a water soluble polymer, for example carboxymethylcellulose, polyvinylpyrrolidone or hydroxypropylmethyl cellulose, preferably carboxymethylcellulose or polyvinylpyrrolidone, can be present.

- 5 Preferably, from 10 to 150mg/g of eletriptan hemisulphate is present.
 Preferably, from 20 to 110mg/g of eletriptan hemisulphate is present.
 Preferably, from 50 to 120mg/g of eletriptan hemisulphate is present.
 Preferably, from 10 to 30% weight/weight of the sulphobutylether-beta-cyclodextrin is present.

- 10 Preferably, from 15 to 25% weight/weight of the sulphobutylether-beta-cyclodextrin is present.

 Preferably, from 0.10 to 1.0% weight/weight of ascorbic acid is present.
 Preferably, from 0.25 to 0.80% weight/weight of ascorbic acid is present.
 Preferably, from 0.30 to 0.60% weight/weight of ascorbic acid is present.

- 15 Preferably, from 5.0 to 30.0% weight/weight of glycerol is present.
 Preferably, from 10.0 to 25.0% weight/weight of glycerol is present.
 Preferably, from 10.0 to 20.0% weight/weight of glycerol is present.
 Preferably, from 0.05 to 0.5% weight/weight of carboxymethylcellulose or polyvinylpyrrolidone is present.

- 20 Preferably, from 0.05 to 0.2%, most preferably, from 0.1 to 0.2%, weight/weight of carboxymethylcellulose or polyvinylpyrrolidone is present.

 Preferably, an aqueous pharmaceutical formulation is adjusted to a pH of from 4.0 to 9.0.

- 25 Preferably, an aqueous pharmaceutical formulation is adjusted to a pH of from 4.0 to 7.0.

 Preferably, an aqueous pharmaceutical formulation is adjusted to a pH of from 4.0 to 5.0.

 The following Examples illustrate the preparation of aqueous pharmaceutical formulations including the complex of the present invention.

- 30 The formulations were prepared by the addition of eletriptan hemisulphate, the sulphobutylether-beta-cyclodextrin, glycerol or polyvinyl pyrrolidone, and

ascorbic acid, to water (sufficient quantity to represent 80% of the final volume of the required formulation). The mixture was stirred to dissolve the solids and the resulting solution adjusted to the required pH using 1M aqueous sodium hydroxide solution. Water was then added to achieve the final volume required.

- 5 The sulphobutylether-beta-cyclodextrin used has an average sulphobutylether substitution of 6.5 per cyclodextrin molecule, and each sulphobutylether unit was present as its sodium salt.

EXAMPLE 1

A formulation suitable for intranasal administration is an aqueous formulation comprising:

- 5 100mg/g of eletriptan hemisulphate;
20% weight/weight of the sulphobutylether-beta-cyclodextrin;
15% weight/weight of glycerol; and
0.5% weight/weight of ascorbic acid:
with the formulation adjusted to from pH 4.0 to 5.0, preferably about pH 4.5,
- 10 using aqueous sodium hydroxide solution.

EXAMPLE 2

A formulation suitable for intranasal administration is an aqueous formulation

- 15 comprising:
80mg/g of eletriptan hemisulphate;
20% weight/weight of the sulphobutylether-beta-cyclodextrin;
0.15% weight/weight of polyvinylpyrrolidone; and
0.5% weight /weight ascorbic acid:
- 20 with the composition adjusted to from pH 4.0 to 5.0, preferably about pH 4.5,
using aqueous sodium hydroxide solution.

EXAMPLE 3

- 25 A formulation suitable for intranasal administration is an aqueous formulation comprising:

- 80mg/g of eletriptan hemisulphate;
20% weight/weight of the sulphobutylether-beta-cyclodextrin;
20% weight/weight of glycerol; and
- 30 0.7% weight/weight of ascorbic acid:

with the formulation adjusted to from pH 4.0 to 5.0, preferably about pH 4.5, using aqueous sodium hydroxide solution.

EXAMPLE 4

5

A formulation suitable for intranasal administration is an aqueous formulation comprising:

80mg/g of eletriptan hemisulphate;

20% weight/weight of the sulphobutylether-beta-cyclodextrin;

10 0.10% weight/weight of polyvinylpyrrolidone; and

0.7% weight /weight ascorbic acid:

with the composition adjusted to from pH 4.0 to 5.0, preferably about pH 4.5, using aqueous sodium hydroxide solution.

15

BIOLOGICAL ACTIVITY

The complex of the present invention may be tested for irritant effects on the nasal mucosae by the following method.

5 Test solutions were prepared as follows:

(1) an aqueous solution containing 10% wt/wt hydroxypropyl-beta-cyclodextrin (containing an average number of hydroxypropyl groups per molecule of cyclodextrin of 0.6);

(2) an aqueous solution containing 17% wt/wt of the present
10 sulphobutylether-beta-cyclodextrin;

(3) an aqueous solution containing 10% wt/wt hydroxypropyl-beta-cyclodextrin (containing an average number of hydroxypropyl groups per molecule of cyclodextrin of 0.6) and 50mg/ml of eletriptan hemisulphate; and

(4) an aqueous solution containing 17% wt/wt of the present
15 sulphobutylether-beta-cyclodextrin and 50mg/ml of eletriptan hemisulphate.

All the above solutions were adjusted to pH 4.2 +/- 0.2.

Female Sprague-Dawley rats were used for the studies. The animals were about 7 weeks old with a mean body weight of 199g. Each study was performed on a group of five rats.

20 A 20 microlitre volume of the test solution was instilled into the left nostril of each rat in the study group. The rats in each group were treated once daily with the same test solution for 7 days. The rats in each study group were observed daily for mortality and clinical signs. They were weighed on study days -2, 1 and 7. On study day 8 the rats were sacrificed and macroscopic and
25 histopathological examinations were carried out on the tissues of the respiratory tract only of each rat.

No deaths were recorded. The results showed that test solutions (1), (2) and (4) did not produce any lesions. Test solution (3) induced minimal to mild hyperplasia and metaplasia of the respiratory epithelium and luminal exudate.


30 These findings were interpreted as signs of minimal to mild irritation of the nasal turbinates.

It was concluded that test solution (4), containing a complex of eletriptan hemisulphate and the present sulphobutylether-beta-cyclodextrin, was non-irritant on the nasal mucosae when administered by the intranasal route.

CLAIMS

1. A complex of eletriptan and a cyclodextrin derivative of formula (I):-

5



wherein

R^{1a-g} , R^{2a-g} and R^{3a-g} each independently represent $-OH$ or $-O(CH_2)_4SO_3H$;

10

provided that at least one of R^{1a-g} represents $-O(CH_2)_4SO_3H$;

or a pharmaceutically acceptable salt thereof.

2. A complex as claimed in claim 1, wherein the average number of $-O(CH_2)_4SO_3H$ groups per molecule of the derivative of the formula (I) is in the range of from 6.1 to 6.9.

15

3. A complex as claimed in claim 1 or claim 2 wherein each $-O(CH_2)_4SO_3H$ group present in the derivative of the formula (I) is in the form of an alkali metal salt.

20

4. A complex as claimed in any one of the preceding claims wherein the molar ratio of eletriptan:cyclodextrin derivative of the formula (I) is from 1:1 to 15:1.

5. A complex as claimed in claim 4 wherein the molar ratio of
eletriptan:cyclodextrin derivative of the formula (I) is from 1:1 to 10:1.
6. A complex as claimed in any one of the preceding claims wherein eletriptan
5 is present in the form of the hemisulphate salt.
7. A pharmaceutical formulation including a complex as claimed in any one of
the preceding claims and a pharmaceutically acceptable excipient, diluent or
carrier.
- 10 8. A formulation as claimed in claim 7 wherein from 50 to 120mg/g of eletriptan
hemisulphate is present.
9. A formulation as claimed in claim 7 or 8 wherein from 15 to 25%
15 weight/weight of the sulphobutylether-beta-cyclodextrin is present.
10. A formulation as claimed in claim 7, 8 or 9 including one or more of an anti-
oxidant, a co-solvent and an organic polymer.
- 20 11. A formulation as claimed in claim 10 wherein the anti-oxidant is ascorbic
acid.
12. A formulation as claimed in claim 11 wherein from 0.25 to 0.80%
weight/weight of ascorbic acid is present.
- 25 13. A formulation as claimed in any one of claims 10 to 12 wherein the co-
solvent is glycerol.
14. A formulation as claimed in claim 13 wherein from 10.0 to 25.0%
30 weight/weight of glycerol is present.

15. A formulation as claimed in any one of claims 10 to 14 wherein the organic polymer is carboxymethylcellulose or polyvinylpyrrolidone.
16. A formulation as claimed in claim 15 wherein from 0.05 to 0.20%
5 weight/weight of carboxymethylcellulose or polyvinylpyrrolidone is present.
17. A formulation as claimed in any one of claims 7 to 16 that is in the form of an aqueous solution.
- 10 18. An aqueous formulation as claimed in claim 17 that has a pH of from 4.0 to 5.0.
19. A formulation as claimed in any one of claims 7 to 18 which is adapted for parenteral administration.
- 15 20. A formulation as claimed in any one of claims 7 to 18 which is adapted for intranasal administration.
21. A formulation as claimed in any one of claims 7 to 18 which is adapted for
20 inhalation.
22. A formulation as claimed in claim 7 that is an aqueous solution comprising:
80mg/g of eletriptan hemisulphate;
25 20% weight/weight of the sulphobutylether-beta-cyclodextrin as defined in claim 1 having an average sulphobutylether substitution of 6.5 per cyclodextrin molecule with each sulphobutylether unit present as its sodium salt;
20% weight/weight of glycerol; and
30 0.7% weight/weight of ascorbic acid:

with the formulation having been adjusted to from pH 4.0 to 5.0, preferably about pH 4.5, using aqueous sodium hydroxide solution.

23. A formulation as claimed in claim 7 that is an aqueous solution comprising:

5 80mg/g of eletriptan hemisulphate;
20% weight/weight of the sulphobutylether-beta-cyclodextrin as defined in claim 1 having an average sulphobutylether substitution of 6.5 per cyclodextrin molecule with each sulphobutylether unit present as its sodium salt;

10 0.10% weight/weight of polyvinylpyrrolidone; and
0.7% weight /weight ascorbic acid:
with the composition having been adjusted to from pH 4.0 to 5.0, preferably about pH 4.5, using aqueous sodium hydroxide solution.

15 24. A complex as claimed in any one of claims 1 to 6 or a pharmaceutically acceptable formulation thereof as claimed in any one of claims 7 to 23 for use as a medicament.

20 25. The use of a complex as claimed in any one of claims 1 to 6 or of a pharmaceutically acceptable formulation thereof as claimed in any one of claims 7 to 23 for the manufacture of a medicament for the treatment of a disease for which a 5HT_{1B/1D} receptor agonist is indicated.

25 26. The use of a complex as claimed in any one of claims 1 to 6 or a pharmaceutically acceptable formulation thereof as claimed in any one of claims 7 to 23 for the manufacture of a medicament for the treatment of migraine or the prevention of migraine recurrence.

30 27. A method of treatment of a mammal to treat a disease for which a 5HT_{1B/1D} receptor agonist is indicated including treating said mammal with an effective amount of a complex as claimed in any one of claims 1 to 6 or a

pharmaceutically acceptable formulation thereof as claimed in any one of claims 7 to 23.

28. A method of treatment of a mammal to treat migraine or to prevent migraine
5 recurrence including treating said mammal with an effective amount of a complex as claimed in any one of claims 1 to 6 or a pharmaceutically acceptable formulation thereof as claimed in any one of claims 7 to 23.
29. A process for the preparation of a complex as defined in claim 1 which
10 comprises combining eletriptan, or a pharmaceutically acceptable salt thereof, with the cyclodextrin derivative, or a pharmaceutically acceptable salt thereof.
30. A process for the preparation of a formulation as claimed in claim 7 which
15 comprises combining either (i) a complex as defined in claim 1, or (ii) eletriptan, or a pharmaceutically acceptable salt thereof, and the cyclodextrin derivative, or a pharmaceutically acceptable salt thereof, with a pharmaceutically acceptable excipient, diluent or carrier.

INTERNATIONAL SEARCH REPORT

Int. tional Application No
PCT/IB 00/00746

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K47/48

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, EMBASE, BIOSIS, EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category ²	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 91 11172 A (UNIV KANSAS) 8 August 1991 (1991-08-08) cited in the application claims	1-30
Y	WO 94 02518 A (UNIV KANSAS) 3 February 1994 (1994-02-03) cited in the application claims 1,30	1-30
P,Y	WO 00 06161 A (JACKSON NEVILLE COLIN ; PFIZER LTD (GB); UDEN STEPHEN (GB); PFIZER) 10 February 2000 (2000-02-10) cited in the application claims 1,8,10	1-30

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

² Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

20 September 2000

Date of mailing of the international search report

28/09/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Berte, M

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 00/00746

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9111172 A	08-08-1991	US 5134127 A AT 170742 T AU 646020 B AU 7236491 A CA 2074186 A DE 69130165 D DE 69130165 T EP 0512050 A JP 2722277 B JP 5504783 T KR 166088 B RU 2099354 C US 5376645 A	28-07-1992 15-09-1998 03-02-1994 21-08-1991 24-07-1991 15-10-1998 29-04-1999 11-11-1992 04-03-1998 22-07-1993 15-01-1999 20-12-1997 27-12-1994
WO 9402518 A	03-02-1994	US 5376645 A AU 672814 B AU 4779993 A CA 2119154 A,C EP 0620828 A JP 6511513 T RU 2113442 C	27-12-1994 17-10-1996 14-02-1994 03-02-1994 26-10-1994 22-12-1994 20-06-1998
WO 0006161 A	10-02-2000	AU 3952199 A	21-02-2000